

Daporex[®]

Dapoxetine Film coated Tablet

Presentation:

Daporex[®] 30 Tablet: Each film coated tablet contains Dapoxetine Hydrochloride INN equivalent to Dapoxetine 30 mg.

Daporex[®] 60 Tablet: Each film coated tablet contains Dapoxetine Hydrochloride INN equivalent to Dapoxetine 60 mg.

Description:

Dapoxetine Hydrochloride is a white powder substance and watersoluble. Dapoxetine Hydrochloride works by inhibiting the serotonin transporter, increasing serotonin's action at the post synaptic cleft and as a consequence promoting ejaculatory delay. Dapoxetine Hydrochloride is absorbed and eliminated rapidly from the body. Its fast acting property makes it suitable for the treatment of Premature Ejaculation. Dapoxetine Hydrochloride is distributed rapidly in the body. Greater than 99% of Dapoxetine Hydrochloride is bound to the plasma protein. The mean steady state volume is 162 L. Its initial half-life is 1.31hours. Dapoxetine Hydrochloride is metabolized extensively in the liver and kidney by multiple enzymes such as CYP2D6, CYP3A4, and flavin mono-oxygenase 1 (FMO1).

Indications:

Daporex[®] is indicated for the treatment of -

- Premature ejaculation.
- Poor control over ejaculation.

Dosage and administration:

The recommended starting dose for all patients is 30 mg, taken as needed approximately 30-60 minutes prior to sexual activity. Dose may be increased to the maximum recommended of 60 mg.

Contraindications:

Dapoxetine Hydrochloride is contraindicated to whom who has hypersensitivity to it or any component of this product.

Precaution:

Dapoxetine Hydrochloride is only indicated in men with premature ejaculation.

Drug Interactions:

Potential serious reactions with MAOIs. Inhibits metabolism of thioridazine. Risk of serotonergic-associated effects with serotonergic medicinal/herbal products (including L-tryptophan, triptans, tramadol, linezolid, SSRIs, SNRIs, lithium & St. John's wort prep). CNS-active medicinal products. Possible reduction of clearance with CYP2D6, CYP3A4 & flavin monooxygenase 1 inhibitors.

Side Effects:

Side effects are classified as low compared with many drugs. Such as dizziness, nausea, dry mouth, flatulence etc.

Storage Condition:

Store in a cool and dry place, away from light. Keep out of the reach of children.

Packaging:

Daporex[®] 30 Tablet: Each box contains 2x4 tablets in Alu-Alu blister pack.

Daporex[®] 60 Tablet: Each box contains 1x4 tablets in Alu-Alu blister pack.



Manufactured by
EURO Pharma Ltd.
In Pursuit of Excellence
Dhaka, Bangladesh

® Regd. Trademark